

Department of Biotechnology

Atmiya University



By : Niva Dudani

Sem: M.sc. 4th semester

Enrollment no.:210621014

Place : Atmiya University

Guided by: Dr.Preetam Joshi

Topic : Study on compatibility of natural IAA isolation from bacteria (a potential) in tissue culture growth *Bryophyllum* plant

Study on compatibility of natural IAA isolation from bacteria (a potential) in tissue culture growth *Bryophyllum* plant

Abstract

Plant tissue culture is a method used to grow and propagate plants outside of their native habitat in a controlled laboratory setting. The purpose of the current investigation was to clarify the potential function of bacterial auxin in plant tissue culture. In order to achieve this, MS basal medium was supplemented with bacterial supernatant containing auxin, and the effect of this addition on in vitro plant growth was investigated. With internodal explants inoculated on MS + BS, callus induction was observed at the apex of explant at 1 and 2 ml, where adventitious root induction is observed in ND1, ND2 and ND-3. In treatment plant ND3 shows significant increase of carbohydrate and root length with number of advantage root number was increase as compared to other ND1. In comparison of control plant no one is increase root num or length, were not reported increase of in terms of biomass carbohydrates and protein.

Keywords

Bryophyllum, *in-vitro* culture, Bacterial auxin , Somatic embryogenesis

INTRODUCTION

Bryophyllum may be a herb which develop broadly in Africa , tropical America, China , India , and Australia. This herb is contains a wide extend of alkaloids, triterpenes, glycosides, flavonoids, steroids, lipids and natural acids. The plant is broadly utilized in conventional pharmaceutical for the treatment of assortment of sicknesses, mending properties are moreover fundamental strength. *Bryophyllum* is develops all over India sodden and warm ranges. The plant, *Bryophyllum* is acknowledged as a home grown cure in a few parts of the world. Numerous of the *Bryophyllum's* conventional employments have been approved by creature ponders and clinical examinations on plant. Numerous of the *Bryophyllum's* conventional employments have been affirmed by the clinical investigate conducted so distant on the plant. The conventional utilize for irresistible conditions is bolstered by inquire about demonstrating that *Bryophyllum* takes off have antibacterial, antiviral and antifungal movement. The leaf and leaf juice have illustrated critical antibacterial movement towards Staphylococcus, Escherichia coli, Shigella, Bacillus and Pseudomonas, counting a few strains of multi-drug safe microscopic organisms.

Succulent plants in the genus *Bryophyllum* are frequently employed in plant tissue culture because of their capacity to regenerate entire plants from tiny explants or callus tissues. Somatic embryogenesis is the term for this process. Undifferentiated cells are propelled into the formation of embryos during somatic embryogenesis(1), which can later grow into whole plants. *Bryophyllum* is especially helpful in this process because it is simple to regenerate and may quickly produce a large number of plants. Moreover, *Bryophyllum* can be used in plant tissue culture to examine how different growth regulators affect the growth and development of plants. Due to its therapeutic benefits, *Bryophyllum* has been utilised in traditional medicine, particularly for the treatment of infections and wounds (2, 3).

Plant tissue culture is a method used to grow and propagate plants outside of their native habitat in a controlled laboratory setting. The growth medium used to assist plant development and growth is one of the essential elements in plant tissue culture. The Murashige and Skoog (MS) medium, created in 1962 by plant scientists Toshio Murashige and Folke Skoog, is one of the most frequently used growth media(4,5).

In order to produce a balanced and optimal nutrient environment for plant tissue culture, the MS medium is a sophisticated blend of salts, vitamins, and plant growth regulators(6,7). It has the ideal ratios of macronutrients like nitrogen, phosphorous, and potassium, as well as micronutrients like iron, manganese, and zinc, to support the growth of plants(8). A variety of vitamins, including thiamine, pyridoxine, and nicotinic acid, which are necessary for plant growth and development, are also included in the medium.

The primary endogenous regulators of numerous aspects of plant growth and development are plant hormones. Auxin, one of the most thoroughly investigated hormones, controls In plants, cells divide, lengthen, differentiate, and form patterns (9).Both naturally occurring chemicals and closely related manufactured compounds with comparable actions are classified as auxins. The hormones auxin and cytokinin regulate the growth of plants through a variety of intricate interactions. Auxin-cytokinin ratio influences the development of lateral roots and the growth of shoot axillary buds (10). (11).

Auxin can also be biosynthesized in lower plants(12). IAA that is physiologically active and produced by organisms like bacteria may have significant effects on plant growth and development. According to (13), over 80% of bacteria isolated from plant rhizospheres may produce indole-3-acetic acid. Because its addition to bacterial cultures that produce IAAs encourages and boosts IAA synthesis, L-tryptophan is also regarded as the IAA precursor in bacteria, just like it is in plants (13,14). For the rhizosphere microflora, root exudates constitute a natural supply of L-tryptophan, which may improve auxin biosynthesis in the rhizosphere (15,16).

The purpose of the current investigation was to clarify the potential function of bacterial auxin in plant tissue culture. In order to achieve this, MS basal medium was supplemented with bacterial supernatant

containing auxin, and the effect of this addition on *in vitro* plant growth was investigated. Think about have illustrated the potential of rhizobacteria to deliver IAA and their related useful impacts when presented into *in vitro* micropropagation conventions at the establishing arrange. Plant development advancing microbes (PGPB) have been appeared to be able of colonizing the rhizosphere these straightforwardly and in a roundabout way impact the development of the has. The utilize of bacteria in plant development tests has in this manner expanded significantly within the final few decades. The utilize of bacteria to advance plant development beneath dry spell conditions. In any case, the utilize of bacteria specifically in *in vitro* societies is still in its earliest stages: more commonly IAA is produced from bacteria and after that connected discretely. In this manner, the point of this think about was to evaluate the use of IAA from natural sources within the *in vitro* rhizogenesis and different growth stages in *bryophyllum*.

Material Method

Explant preparation and culture establishment: *Bryophyllum* leaf cultures were created using the method (17). Plants in good health were purchased at the botanical garden. Fresh leaves were chosen, trimmed, and rinsed with tap water to remove dust before going through two detergent washes and being sterilised by soaking them in 70% ethanol for one minute. By treating the explant with 1% sodium hypochlorite for 10 minutes and then rinsing it three to four times in sterile distilled water, the explant was further sterilised. The standard Murashige and Skoog's (1962) medium, which contains 3.0 mg L⁻¹ BAP, 0.01 mg L⁻¹ NAA, 0.8% agar, and 3.0% sucrose, was used for aseptically inoculating the explant. Every three weeks after the initial establishment of cultures, routine sub-culturing was carried out. Cultures were maintained

Experiment design: The MS medium (Murashige and Skoog, 1962) was the base medium utilised in this investigation. It contained 8 g agar l⁻¹, 30 g sucrose l⁻¹, 100 mg myoinositol l⁻¹, and 0.4 mg thiamine-HCl l⁻¹ as carbon sources. Each test tube (2.5 15 cm) received five millilitres of MS medium, which was utilised in various combinations. Filter-sterilized bacterial supernatants were added to three groups of combinations in varying amounts—1, 2, and 5 ml each in 50 ml of MS media. Three different MS medium combinations were used: DS1: MS + BS(1 ml), DS2: MS + BS(2 ml), DS3: MS + BS(5 ml), and DS4: MS + IAA(2 mg).(18)

Measurement of growth parameters: Average root length, total root number and total biomass in terms of fresh and dry weight were measured. For measurement of biomass (fresh weight and dry weight), propagules obtained from each treatment were taken out and the fresh weight was measured using an electronic top pan balance. For dry weight calculation, after measuring the fresh weight those fresh shoots were kept in an oven at 62°C for drying till constant weight.

Chlorophyll contents: The chlorophyll contents were calculated as per the method described by (19). 500 mg of shoots were weighed and ground in pestle and mortar with 80 % acetone under dark conditions. Extracts were centrifuged at 10, 000 rpm and the supernatant was used to measure absorbance on spectrophotometer (UV-1800 Shimadzu, Japan).

Total phenols: The total phenol content was measured as per the method described (20) 500 mg of shoots were weighed and crushed in pestle and mortar in 70 % methanol. The extracts were centrifuged at 10,000 rpm for 15 minutes. The clear supernatants were used for quantitative determination of total phenol content. For each reaction 500 µL methanolic extract was taken in a test tube and to this 1.0 mL suitably diluted (1:1 ratio of reagent and DDW) Folin Ciocalteu's reagent was added followed by 2.0 mL of Na₂CO₃ (20 % w/v) solution. The test tubes were heated in boiling water bath with intermittent shaking for about 1.0 min. Tubes were subsequently cooled under running tap water. The blue colored product was diluted to 25 mL by adding DDW and the percent transmittance was measured at 650 nm in a spectrophotometer (UV-1800 Shimadzu, Japan). The total phenol concentration in each sample was estimated with the help of standard curve prepared using different concentrations (10-100 µg) of caffeic acid.

Total carbohydrates: Quantitative estimation of total carbohydrate content was carried out as per method described by Tandon (21). In vitro derived propagules were homogenized in 0.1 M phosphate buffer (pH 7.0) and the homogenates were centrifuged at 10,000 rpm for 15 min. For each reaction 15 µL of supernatant was mixed with 4.0 mL of 0.2 % Anthrone reagent (in conc. H₂SO₄) and placed in boiling water bath for five minutes. The absorbance was recorded at 610 nm wave length. The total carbohydrate contents were determined using standard curve prepared from various concentrations of glucose.

Total protein: Quantitative estimation of total protein was performed as per (22) One mL of the suitably diluted crude tissue extract (the supernatant) was mixed with 5.0 mL of Coomassie Brilliant Blue G-250 dye (Bradford reagent) and transmittance of the resultant solution (coloured complex) was read with spectrophotometer (UV-1800 Shimadzu, Japan) at 595 nm. The amount of protein was determined using standard curve prepared using various concentrations of albumin protein.

Table 1 IAA Composition Different Concentration

Media	PGRs	Isolated PGRs
ND4 (Control)	IAA-2mg/L	-
ND1	-	20 ml/L
ND2	-	40 ml/L
ND3	-	100 ml/L

Table 2: Ingredient of MS Media

Ingredients	Quantity Required or 1000 ml

Stock A	50 ML
Stock B	5 ML
Stock C	5 ML
Stock D	5 ML
Agar Agar	0.8%
Sucrose	0.3%
BAP	2 mg
IAA	0.75 mg

Result and discussion

With internodal explants inoculated on MS + BS, callus induction was observed at the apex of explants at 1 and 2 ml as compared to 5 ml, where adventitious root induction was observed in DS1, DS2 and DS3. In MS medium supplemented with standard IAA adventitious root induction was observed on all treatments with maximum induction as compared to treatment plant. whereas in MS + IAA can also use for shoot induction as IAA alone can

Table no.3 Effect of bacterial Auxin on In vitro plant root growth and biochemical parameters.

Media	Fresh Weight	Dry Weight	Root Length	Root Number	Total Carbohydrate	Total protein
ND1	8 gm	1.2 gm	2 cm	18	100 mg g ⁻¹	68 mg g ⁻¹
ND2	8.5 gm	1.5 gm	2 cm	20	110 mg g ⁻¹	79 mg g ⁻¹
ND3	10 gm	1.9 gm	5 Cm	25	125 mg g ⁻¹	75 mg g ⁻¹

ND4 (Control)	12 gm	2 gm	3 cm	28	130 mg g ⁻¹	85 mg g ⁻¹
------------------	-------	------	------	----	------------------------	-----------------------

induce callus on leaf segments, and *in vitro* experiments have demonstrated that coconut water in combination with synthetic auxins can be used for shoot induction and multiplication (Cui et al. 2004; Loc et al. 2005). In term of biochemical total carbohydrate was higher found in control plant as compared to treatment plant. In treatment plant ND3 shows significant increase of carbohydrate and root length with number of advantage root number was increase as compare to other ND1 and ND2 but in compare of control plant no one is increase root num or length , were not reported increase of in term of biomass carbohydrate an protein.(table 3)



Fig :1 Effect of Bacterial Auxin on *In vitro* grown plant root growth.




Fig:2 Adventitious root induction from *In vitro* plant growth

References

1. Chen, J. T., & Chang, W. C. (2002). Effects of tissue culture conditions and explant characteristics on direct somatic embryogenesis in *Oncidium Gower Ramsey*'. *Plant cell, tissue and organ culture*, 69, 41-44.
2. Al-Snafi, A. E. (2013). The Chemical constituents and pharmacological effects of *Bryophyllum calycinum*. A review. *Journal of Pharma Sciences and Research*, 4(12), 171-176.
3. Berleth T, Sachs T (2001) Plant morphogenesis: long-distance coordination and local patterning. *Curr Opin Plant Biol* 4:57–62
4. Gamborg, O. L., Murashige, T., Thorpe, T. A., & Vasil, I. K. (1976). Plant tissue culture media. *In vitro*, 12(7), 473-478.
5. George, E. F., Hall, M. A., & Klerk, G. J. D. (2008). The components of plant tissue culture media I: macro-and micro-nutrients. *Plant Propagation by Tissue Culture: Volume 1. The Background*, 65-113.
6. Munthali, C., Kinoshita, R., Onishi, K., Rakotondrafara, A., Mikami, K., Koike, M., ... & Aiuchi, D. (2022). A Model Nutrition Control System in Potato Tissue Culture and Its Influence on Plant Elemental Composition. *Plants*, 11(20), 2718.
7. Saad, A. I., & Elshahed, A. M. (2012). Plant tissue culture media. *Recent advances in plant in vitro culture*, 30-40.
8. Cline MG (1996) Exogenous auxin effects on lateral bud outgrowth in decapitated shoots. *Ann Bot* 78:255–266
9. Ehlert, B., Schöttler, M. A., Tischendorf, G., Ludwig-Müller, J., & Bock, R. (2008). The paramutated SULFUREA locus of tomato is involved in auxin biosynthesis. *Journal of Experimental Botany*, 59(13), 3635-3647.
10. Costacurta A, Vanderleyden J (1995) Synthesis of phytohormones by plant-associated bacteria. *Crit Rev Microbiol* 21:1–18
11. Park M, Kim C, Yang J, Lee H, Shin W, Kim S, Sa T (2005) Isolation and characterization of diazotrophic growth promoting bacteria from rhizosphere of agricultural crops of Korea. *Microbiol Res* 160:127–133
12. Tsavkelova EA, Cherdyntseva TA, Netrusov AI (2005) Auxin production by bacteria associated with orchid roots. *Microbiology* 74:46–53
13. Kravchenko LV, Borovkov AV, Pshikvil Z (1991) The possibility of auxin biosynthesis in wheat rhizosphere by associated nitrogen-fixing bacteria. *Microbiology* 60:927–931

14. Martens DA, Frankenberger WT Jr (1994) Assimilation of exogenous 2-14C-indole acetic acid and 3-14C-tryptophan exposed to the roots of three wheat varieties. *Plant Soil* 166:281–290
15. Patten CL, Glick BR (2002) Role of *Pseudomonas putida* indoleacetic acid in development of host plant root system. *Appl Environ Microbiol* 68:3795–3801
16. Souza, J. A., Bettoni, J. C., Dalla Costa, M., Baldissera, T. C., dos Passos, J. F. M., & Primieri, S. (2022). 3-IAA on in vitro rooting and acclimatization of apple rootstock. *Communications*, 12, 016-023.
17. Palta, J. P. (1990). Leaf chlorophyll content. *Remote sensing reviews*, 5(1), 207-213.
18. Mathur, N., & Vyas, A. (1996). Biochemical changes in *Ziziphus xylopyrus* by VA mycorrhizae. *Botanical Bulletin of Academia Sinica*, 37.
19. Mathur, N., & Vyas, A. (1996). Biochemical changes in *Ziziphus xylopyrus* by VA mycorrhizae. *Botanical Bulletin of Academia Sinica*, 37.
20. Bradford, M. M. (1976). A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Analytical biochemistry*, 72(1-2), 248-254.