Accebte

Dual Role of Par-4 in abrogation of EMT and switching on Mesenchymal to

Epithelial Transition (MET) in metastatic pancreatic cancer cells[†]

Archana Katoch^{a, b}, Sujit Suklabaidya^c, Souneek Chakraborty^{a, b}, Debasis Nayak^{a, b},

Reyaz Ur Rasool^{a, b}, Deepak Sharma^d, Debaraj Mukherjee^d, Mir Mohd Faheem^b, Anmol

Kumar^e, Parduman Raj Sharma^b, Shantibhusan Senapati^c, Lekha Dinesh Kumar^e, and

Anindya Goswami ^{a, b, *}

^aAcademy of Scientific & Innovative Research (AcSIR), New Delhi, India.

^bCancer Pharmacology Division, Indian Institute of Integrative Medicine (CSIR), Canal Road, Jammu, Jammu and Kashmir - 180001, India.

^cTumor Microenvironment and Animal Models Lab, Institute of Life Sciences (ILS), Nalco Square Bhubaneswar, Orissa-751023, India

^dNatural Product Chemistry, Indian Institute of Integrative Medicine (CSIR), Canal Road, Jammu, Jammu and Kashmir - 180001, India

^eCancer Biology Division, Center for Cellular and Molecular Biology (CCMB), Uppal Road, Hyderabad, Telangana - 500007, India

[†]This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: [10.1002/mc.22828]

Additional Supporting Information may be found in the online version of this article.

Received 24 November 2017; Revised 10 April 2018; Accepted 17 April 2018 Molecular Carcinogenesis This article is protected by copyright. All rights reserved DOI 10.1002/mc.22828