

Contents

Title	Content	Page No.
Declaration by the Candidate		I
Certificate by Supervisor		III
Acknowledgement		V
Contents		IX
List of Tables		XIII
List of Figures		XVII
List of Abbreviation		XXI-
Abstract		XXV
Chapter 1	Introduction	1

1.1 Introduction

1.1.1 Research region

1.1.2 Background Significance

1.2 PGPR

1.2.1 Optimal PGPR

1.2.2 Role of PGPR in the enhancement of Plant Growth

1.2.3 PGPR and Plant Hormones

1.2.4 Nutrient availability for Plant Growth

1.2.5 Enzymes by PGPR

1.2.6 Abiotic Stress Tolerance in Plants

1.3 Macronutrients and Micronutrients

1.3.1 Nanotechnology and PGPR

1.3.2 Other nanomaterials

1.4 Research Objectives

1.5 Implications

Chapter 2	Review Literature	17
------------------	--------------------------	-----------

2.1 Plant Growth Promoting Rhizobacteria (PGPR)

2.1.1 Diversity of PGPR

2.1.2 The Rhizosphere: A Hotspot for PGPR Activity

2.1.3 Key Roles of PGPR

2.1.4 Importance of PGPR in Agriculture

2.1.5 Mechanisms of Action

2.1.6 Role of *Bacillus* and *Pseudomonas* species as a plant growth promoter in

-
- enhancement of crop production
 - 2.2 PGPR in the enhancement of crop production
 - 2.3 Metal oxide nanoparticles in agriculture
 - 2.3.1 Metal oxide nanoparticles with PGPR
 - 2.4 Co-application of ZnO NPs and PGPR in plants

Chapter 3 Materials and Methods

35

- 3.1 Isolation and screening of plant growth promoting rhizobacteria strain
 - 3.1.1 Collection of Sample and Isolation of Rhizobacteria
 - 3.1.2 Primary Isolation
- 3.2 Characterization of the Isolates for PGP traits
 - 3.2.1 Indole Acetic Acid Production
 - 3.2.2 Ammonia Production
 - 3.2.3 Nitrogen Fixation
 - 3.2.4 Zinc Solubilization
 - 3.2.5 Potassium Solubilization
 - 3.2.6 Chitin Hydrolysis
 - 3.2.7 Hydrogen Cyanide Production
 - 3.2.8 Phosphate Solubilization Test
 - 3.2.9 Gibberellin (GA) production
 - 3.2.10 Siderophore production
- 3.3 Identification of Potent PGPR
 - 3.3.1 Gram staining and KOH method
 - 3.3.2 Molecular identification of PGPR isolate by 16s rRNA sequencing
- 3.4 Synthesis of ZnO NPs using chemical method
 - 3.4.1 Characterization of Synthesized ZnO
- 3.5 Bacterial Growth Curve
- 3.6 Preparation of bacterial culture
- 3.7 Seed Germination and Vigor Index
- 3.8 Optimization of synthesized ZnO NPs on plant growth
 - 3.8.1 Pot Assay
 - 3.8.2 Measurements of Physiochemical and Biochemical Parameters
- 3.9 Optimization of synthesized ZnO NPs on plant growth with 3 potent strains
 - 3.9.1 Physical and Biochemical Parameters (Data Recording and Related Procedures)
 - Effect of different growth condition on plant growth

Chapter 4	Results and Discussion (Isolation and Molecular Identification of the Plant Growth-Promoting Rhizobacteria-PGPR)	51
4.1	Primary and Secondary Screening (PGP Traits)	
4.2	Identification of Potent PGPR	
4.2.1	Gram Staining and KOH (potassium hydroxide) Method	
4.2.2	Molecular Identification of Potent PGPR	
Chapter 5	Results and Discussion (Chemical Synthesis, Characterization and Optimization Study of ZnO NPs)	63
5.1	Chemically Synthesized ZnO NPs	
5.2	Characterization of ZnO NPs	
5.3	Bacterial growth curve	
5.4	Seed priming and vigor index	
5.5	Optimization of the effective concentration of ZnO NPs for plant growth	
5.5.1	Physical parameters	
Chapter 6	Results and Discussion (Pot Experiment on the Co-application of Three Potent PGPRs and Optimized ZnO NP Concentrations)	77
6.1	Effect of PGPR and ZnO NPs on plant growth	
6.1.1	Physical parameters	
6.1.2	Biochemical parameters	
6.1.2.1	Estimation of Total Chlorophyll content for RG8 (<i>Pseudomonas songnenensis</i>)	
6.1.2.2	Estimation of Carotenoid content for RG8 (<i>Pseudomonas songnenensis</i>)	
6.1.2.3	Estimation of Flavonoids content for RG8 (<i>Pseudomonas songnenensis</i>)	
6.1.2.4	Estimation of Proline content for RG8 (<i>Pseudomonas songnenensis</i>)	
6.1.2.5	Estimation of Total Protein content for RG8 (<i>Pseudomonas songnenensis</i>)	
6.1.2.6	Estimation of Sugar content for RG8 (<i>Pseudomonas songnenensis</i>)	
6.1.2.7	Estimation of Total Chlorophyll content for RG12 (<i>Bacillus haynesii</i>)	
6.1.2.8	Estimation of Carotenoid content for RG12 (<i>Bacillus haynesii</i>)	
6.1.2.9	Estimation of Flavonoids content for RG12 (<i>Bacillus haynesii</i>)	
6.1.2.10	Estimation of Proline content for RG12 (<i>Bacillus haynesii</i>)	

-
- 6.1.2.11 Estimation of Total Protein Content for RG12 (*Bacillus haynesii*)
 - 6.1.2.12 Estimation of Sugar content for RG12 (*Bacillus haynesii*)
 - 6.1.2.13 Estimation of Total Chlorophyll content for RGKP3 (*Priestia megaterium*)
 - 6.1.2.14 Estimation of Carotenoid content for RGKP3 (*Priestia megaterium*)
 - 6.1.2.15 Estimation of Flavonoids content for RGKP3 (*Priestia megaterium*)
 - 6.1.2.16 Estimation of Proline content for RGKP3 (*Priestia megaterium*)
 - 6.1.2.17 Estimation of Total Protein Content for RGKP3 (*Priestia megaterium*)
 - 6.1.2.18 Estimation of Sugar content for RGKP3 (*Priestia megaterium*)

Chapter 7	Results and Discussion (Comparative Analysis of Three Potent PGPR Strains in a Pot Experiment: Results with ANOVA Study)	105
7.1	Pot experiment	
7.1.1	Physical parameters	
7.1.2	Biochemical parameters	
7.1.2.1	Total Chlorophyll Content	
7.1.2.2	Carotenoid Content	
7.1.2.3	Flavonoids Content	
7.1.2.4	Proline Content	
7.1.2.5	Total Protein Content	
7.1.2.6	Sugar Content	
Chapter 8	Summary	117
	Bibliography	123
Appendix A	Plagiarism Report	
Appendix B	Publications	

List of Figures

Figure No.	Title	Page No.
Figure 1.1	Schematic representation of Plant Growth by PGPR	5
Figure 1.2	The role and function of various soil minerals, microorganisms, macronutrients, and micronutrients in plant growth and development	10
Figure 4.1	(a) Qualitative analysis of IAA production of isolates; (b) Qualitative analysis of ammonia production of isolates	51
Figure 4.2	Quantification of IAA production of all positive isolates	52
Figure 4.3	Quantification of ammonia production of eighteen positive isolates	52
Figure 4.4	Qualitative analysis of HCN production of all positive isolates	53
Figure 4.5	Quantitative analysis of HCN production of isolates compared to control	54
Figure 4.6	Quantification of HCN production of twenty-six positive isolates	54
Figure 4.7	(a) Quantitative analysis of phosphate solubilization control with positive result of isolate RGKP3; and (b) Quantification of phosphate solubilization	55
Figure 4.8	Quantitative analysis of Gibberellins (GA) production compared to control	55

Figure 4.9	Quantification of Gibberellins (GA) production	56
Figure 4.10	(a) Chitin hydrolysis with a positive result of isolate RGKP3; and (b) Siderophore production with a positive result of isolate RGKP3	57
Figure 4.11	Potassium solubilization of isolate RGKP3 compared to control	57
Figure 4.12	(a) Nitrogen fixation of isolate RGKP3; and (b) Zinc solubilization of isolate RGKP3	58
Figure 4.13	Gram staining images for all three positive isolates; (a) RG8; (b) RG12; and (c) RGKP3	59
Figure 4.14	Phylogenetic tree showing the evolutionary relationship between RG8, a PGPR isolate, and reference strains from the GenBank database	60
Figure 4.15	Phylogenetic tree showing the evolutionary relationship between RG12, a PGPR isolate, and reference strains from the GenBank database	61
Figure 4.16	Phylogenetic tree showing the evolutionary relationship between RGKP3, a PGPR isolate, and reference strains from the GenBank database	62
Figure 5.1	UV visible spectrophotometric analysis of ZnO NPs	63
Figure 5.2	SEM micrograph of ZnO NPs (white dots represent nanoparticles)	64
Figure 5.3	TEM micrograph of ZnO NPs with an average diameter of 50-70 nm	65
Figure 5.4	XRD analysis of chemically synthesized zinc oxide nanoparticles at 400 ppm concentration	66
Figure 5.5	Bacterial growth curve of (a) Only RG8 and RG8 with 400 ppm ZnO NPs; (b) Only RG12 and RG12	67

	with 400 ppm ZnO NPs; and (c) Only RGKP3 and RGKP3 with 400 ppm ZnO NPs	
Figure 5.6	Groundnut seed Germination with different treatments	69
Figure 5.7	Plants with different treatments in pot experiment after 1 month; Physical parameters of plants with combined PGPR and ZnO NPs at various concentration	69
Figure 5.8	Measurements of fresh and dry weight for different treated plants (Bar represents standard error)	70
Figure 5.9	Measurements of root and shoot length for different treated plants (Bar represents standard error)	70
Figure 5.10	Measurements of physical parameters such as Numbers of Leaves, branches, and roots for various concentrations of ZnO NPs in a pot experiment (Bar represents standard error)	71
Figure 5.11	Estimation of total chlorophyll content for all different treated plants (Bar represents standard error)	72
Figure 5.12	Estimation of produced carotenoid content for different treated plants (Bar represents standard error)	73
Figure 5.13	Estimation of produced flavonoid content for different treated plants (Bar represents standard error)	73
Figure 5.14	Estimation of produced proline content for different treated plants (Bar represents standard error)	74
Figure 5.15	Estimation of produced protein content for different treated plants (Bar represents standard error)	75
Figure 5.16	Estimation of produced total sugar content for different treated plants (Bar represents standard error)	76

Figure 6.1	Physical parameters number of leaves of plants with combined PGPR and ZnO NPs, Only PGPRs, Only NPs, zinc salt, and control (untreated) focusing on SSR expression. Duncan's method was used to compare the means at a 1% probability level, with columns sharing the same letters indicating no significant differences	77
Figure 6.2	Physical parameters number of roots of plants with combined PGPR and ZnO NPs, Only PGPRs, Only NPs, zinc salt, and control (untreated) focusing on SSR expression. Duncan's method was used to compare the means at a 1% probability level, with columns sharing the same letters indicating no significant differences	78
Figure 6.3	Physical parameters number of branches of plants with combined PGPR and ZnO NPs, Only PGPRs, Only NPs, zinc salt, and control (untreated) focusing on SSR expression. Duncan's method was used to compare the means at a 1% probability level, with columns sharing the same letters indicating no significant differences	79
Figure 6.4	Physical parameters number of shoots of plants with combined PGPR and ZnO NPs, Only PGPRs, Only NPs, zinc salt, and control (untreated) focusing on SSR expression. Duncan's method was used to compare the means at a 1% probability level, with columns sharing the same letters indicating no significant differences	80
Figure 6.5	Measurement of root length(cm) of plants with combined PGPR and ZnO NPs, Only PGPRs, Only NPs, zinc salt, and control (untreated) focusing on SSR expression. Duncan's method was used to compare the means at a 1% probability level, with columns sharing the same letters indicating no significant differences	81

- Figure 6.6 Measurement of shoot length(cm) of plants with combined PGPR and ZnO NPs, Only PGPRs, Only NPs, zinc salt and control (untreated) focusing on SSR expression. Duncan's method was used to compare the means at a 1% probability level, with columns sharing the same letters indicating no significant differences 81
- Figure 6.7 Measurement of fresh weight(g) of plants with combined PGPR and ZnO NPs, Only PGPRs, Only NPs, zinc salt, and control (untreated) focusing on SSR expression. Duncan's method was used to compare the means at a 1% probability level, with columns sharing the same letters indicating no significant differences 82
- Figure 6.8 Measurement of dry weight(g) of plants with combined PGPR and ZnO NPs, Only PGPRs, Only NPs, zinc salt, and control (untreated) focusing on SSR expression. Duncan's method was used to compare the means at a 1% probability level, with columns sharing the same letters indicating no significant differences 82
- Figure 6.9 Total chlorophyll content estimation for various treatments, including RG8 (*Pseudomonas songnenensis* with 400 ppm ZnO NPs), zinc salt alone, 400 ppm ZnO NPs alone, and untreated plants, focusing on SSR expression. Duncan's method was used to compare the means at a 1% probability level, with columns sharing the same letters indicating no significant differences 84
- Figure 6.10 Carotenoid content estimation for various treatments, including RG8 (*Pseudomonas songnenensis* with 400 ppm ZnO NPs), zinc salt alone, 400 ppm ZnO NPs alone, and untreated plants, focusing on SSR expression. Duncan's method was used to compare the means at a 1% probability level, with columns sharing the same letters indicating no significant differences 85

- Figure 6.11 Flavonoid content estimation for various treatments, including RG8 (*Pseudomonas songnenensis* with 400 ppm ZnO NPs), zinc salt alone, 400 ppm ZnO NPs alone, and untreated plants, focusing on SSR expression. Duncan's method was used to compare the means at a 1% probability level, with columns sharing the same letters indicating no significant differences 86
- Figure 6.12 Proline content estimation for various treatments, including RG8 (*Pseudomonas songnenensis* with 400 ppm ZnO NPs), zinc salt alone, 400 ppm ZnO NPs alone, and untreated plants, focusing on SSR expression. Duncan's method was used to compare the means at a 1% probability level, with columns sharing the same letters indicating no significant differences 87
- Figure 6.13 Total protein content estimation for various treatments, including RG8 (*Pseudomonas songnenensis* with 400 ppm ZnO NPs), zinc salt alone, 400 ppm ZnO NPs alone, and untreated plants, focusing on SSR expression. Duncan's method was used to compare the means at a 1% probability level, with columns sharing the same letters indicating no significant differences 88
- Figure 6.14 Total sugar content estimation for various treatments, including RG8 (*Pseudomonas songnenensis* with 400 ppm ZnO NPs), zinc salt alone, 400 ppm ZnO NPs alone, and untreated plants, focusing on SSR expression. Duncan's method was used to compare the means at a 1% probability level, with columns sharing the same letters indicating no significant differences 89
- Figure 6.15 Total chlorophyll content estimation for various treatments, including RG12 (*Bacillus haynesii* with 400 ppm ZnO NPs), zinc salt alone, 400 ppm ZnO NPs alone, and untreated plants, focusing on SSR expression. Duncan's method was used to compare 90

the means at a 1% probability level, with columns sharing the same letters indicating no significant differences

- Figure 6.16 Carotenoid content estimation for various 91
treatments, including RG12 (*Bacillus haynesii* with
400 ppm ZnO NPs), zinc salt alone, 400 ppm ZnO
NPs alone, and untreated plants, focusing on SSR
expression. Duncan's method was used to compare
the means at a 1% probability level, with columns
sharing the same letters indicating no significant
differences
- Figure 6.17 Flavonoid content estimation for various 92
treatments, including RG12 (*Bacillus haynesii* with
400 ppm ZnO NPs), zinc salt alone, 400 ppm ZnO
NPs alone, and untreated plants, focusing on SSR
expression. Duncan's method was used to compare
the means at a 1% probability level, with columns
sharing the same letters indicating no significant
differences
- Figure 6.18 Proline content estimation for various treatments, 93
including RG12 (*Bacillus haynesii* with 400 ppm
ZnO NPs), zinc salt alone, 400 ppm ZnO NPs alone,
and untreated plants, focusing on SSR expression.
Duncan's method was used to compare the means at
a 1% probability level, with columns sharing the
same letters indicating no significant differences
- Figure 6.19 Total protein content estimation for various 94
treatments, including RG12 (*Bacillus haynesii* with
400 ppm ZnO NPs), zinc salt alone, 400 ppm ZnO
NPs alone, and untreated plants, focusing on SSR
expression. Duncan's method was used to compare
the means at a 1% probability level, with columns
sharing the same letters indicating no significant
differences
- Figure 6.20 Total sugar content estimation for various 95
treatments, including RG12 (*Bacillus haynesii* with
400 ppm ZnO NPs), zinc salt alone, 400 ppm ZnO

NPs alone, and untreated plants, focusing on SSR expression. Duncan's method was used to compare the means at a 1% probability level, with columns sharing the same letters indicating no significant differences

- Figure 6.21 Total chlorophyll content estimation for various treatments, including RGKP3 (*Priestia megaterium* with 400 ppm ZnO NPs), zinc salt alone, 400 ppm ZnO NPs alone, and untreated plants, focusing on SSR expression. Duncan's method was used to compare the means at a 1% probability level, with columns sharing the same letters indicating no significant differences 96
- Figure 6.22 Carotenoid content estimation for various treatments, including RGKP3 (*Priestia megaterium* with 400 ppm ZnO NPs), zinc salt alone, 400 ppm ZnO NPs alone, and untreated plants, focusing on SSR expression. Duncan's method was used to compare the means at a 1% probability level, with columns sharing the same letters indicating no significant differences 97
- Figure 6.23 Flavonoid content estimation for various treatments, including RGKP3 (*Priestia megaterium* with 400 ppm ZnO NPs), zinc salt alone, 400 ppm ZnO NPs alone, and untreated plants, focusing on SSR expression. Duncan's method was used to compare the means at a 1% probability level, with columns sharing the same letters indicating no significant differences 98
- Figure 6.24 Proline content estimation for various treatments, including RGKP3 (*Priestia megaterium* with 400 ppm ZnO NPs), zinc salt alone, 400 ppm ZnO NPs alone, and untreated plants, focusing on SSR expression. Duncan's method was used to compare the means at a 1% probability level, with columns sharing the same letters indicating no significant differences 99

- Figure 6.25 Total protein content estimation for various treatments, including RGKP3 (*Priestia megaterium* with 400 ppm ZnO NPs), zinc salt alone, 400 ppm ZnO NPs alone, and untreated plants, focusing on SSR expression. Duncan's method was used to compare the means at a 1% probability level, with columns sharing the same letters indicating no significant differences 100
- Figure 6.26 Total sugar content estimation for various treatments, including RGKP3 (*Priestia megaterium* with 400 ppm ZnO NPs), zinc salt alone, 400 ppm ZnO NPs alone, and untreated plants, focusing on SSR expression. Duncan's method was used to compare the means at a 1% probability level, with columns sharing the same letters indicating no significant differences 101
- Figure 6.27 Total Chlorophyll and Carotenoid content of all treated plants (Salt=Zinc acetate-400 ppm RG8=*Pseudomonas songnenensis*, RG8+NPs=*Pseudomonas songnenensis* with 400 ppm ZnO NPs, RG12=*Bacillus haynesii*, RG12+NPs=*Bacillus haynesii* with 400 ppm ZnO NPs, RGKP3=*Priestia megaterium*, RGKP3+NPs=*Priestia megaterium* with 400 ppm ZnO NPs) 101
- Figure 6.28 Flavonoid estimation was performed for all treated plants under the following conditions: Salt treatment (Zinc acetate, 400 ppm), RG8 (*Pseudomonas songnenensis*), RG8+NPs (*Pseudomonas songnenensis* with 400 ppm ZnO NPs), RG12 (*Bacillus haynesii*), RG12+NPs (*Bacillus haynesii* with 400 ppm ZnO NPs), RGKP3 (*Priestia megaterium*), and RGKP3+NPs (*Priestia megaterium* with 400 ppm ZnO NPs) 102
- Figure 6.29 Proline estimation was conducted for all treated plants under the following treatments: Salt (Zinc acetate, 400 ppm), RG8 (*Pseudomonas songnenensis*), RG8+NPs (*Pseudomonas* 102

songnenensis with 400 ppm ZnO NPs), RG12 (*Bacillus haynesii*), RG12+NPs (*Bacillus haynesii* with 400 ppm ZnO NPs), RGKP3 (*Priestia megaterium*), and RGKP3+NPs (*Priestia megaterium* with 400 ppm ZnO NPs)

- Figure 6.30 Protein estimation was carried out for all treated plants under the following treatments: Salt (Zinc acetate, 400 ppm), RG8 (*Pseudomonas songnenensis*), RG8+NPs (*Pseudomonas songnenensis* with 400 ppm ZnO NPs), RG12 (*Bacillus haynesii*), RG12+NPs (*Bacillus haynesii* with 400 ppm ZnO NPs), RGKP3 (*Priestia megaterium*), and RGKP3+NPs (*Priestia megaterium* with 400 ppm ZnO NPs) 103
- Figure 6.31 Total sugar content was measured for all treated plants under the following treatments: Salt (Zinc acetate, 400 ppm), RG8 (*Pseudomonas songnenensis*), RG8+NPs (*Pseudomonas songnenensis* with 400 ppm ZnO NPs), RG12 (*Bacillus haynesii*), RG12+NPs (*Bacillus haynesii* with 400 ppm ZnO NPs), RGKP3 (*Priestia megaterium*), and RGKP3+NPs (*Priestia megaterium* with 400 ppm ZnO NPs) 103
- Figure 7.1 Plants with different treatments in pot experiment after 1 month; Physical parameters number of leaves, roots, shoots, and branches focusing on SSR expression. Duncan's method was used to compare the means at a 1% probability level, with columns sharing the same letters indicating no significant differences 107
- Figure 7.2 (a) Physical parameters such as root length and shoot length and (b) Physical parameters such as fresh weight and dry weight of plants with combined PGPR and ZnO NPs, Only PGPRs, Only NPs, zinc salt and control (untreated) focusing on SSR expression. Duncan's method was used to compare the means at a 1% probability level, with 107

columns sharing the same letters indicating no significant differences

- Figure 7.3 Total Chlorophyll Content Estimation in Treated Plants from the Pot Experiment focusing on SSR expression. Duncan's method was used to compare the means at a 1% probability level, with columns sharing the same letters indicating no significant differences 108
- Figure 7.4 Carotenoid Content Estimation in Treated Plants from the Pot Experiment focusing on SSR expression. Duncan's method was used to compare the means at a 1% probability level, with columns sharing the same letters indicating no significant differences 109
- Figure 7.5 Flavonoid Content Estimation in Treated Plants from the Pot Experiment focusing on SSR expression. Duncan's method was used to compare the means at a 1% probability level, with columns sharing the same letters indicating no significant differences 110
- Figure 7.6 Proline Content Estimation in Treated Plants from the Pot Experiment focusing on SSR expression. Duncan's method was used to compare the means at a 1% probability level, with columns sharing the same letters indicating no significant differences 111
- Figure 7.7 Total Sugar Content Estimation in Treated Plants from the Pot Experiment focusing on SSR expression. Duncan's method was used to compare the means at a 1% probability level, with columns sharing the same letters indicating no significant differences 112
- Figure 7.8 Total Protein Content Estimation in Treated Plants from the Pot Experiment focusing on SSR expression. Duncan's method was used to compare the means at a 1% probability level, with columns 113

	sharing the same letters indicating no significant differences	
Figure 7.9	Groundnut Plants with different treatments in pot experiment after 1 month of in vitro experiment with combined application of three potent PGPR (RG8, RG12 and RGKP3) strains with ZnO NPs	114
Figure 8.1	Physical parameters such as No. of roots, No. of branch, No. of leaves, No. of shoots h of Groundnut Plants with different treatments in pot experiment after 1 month of in vitro experiment with combined application of three potent PGPR (RG8, RG12 and RGKP3) strain with ZnO NPs. focusing on SSR expression. Duncan's method was used to compare the means at a 1% probability level, with columns sharing the same letters indicating no significant differences	120
Figure 8.2	Physical parameters such as a) Fresh and Dry weights b) Root and shoot length of Groundnut Plants with different treatments in pot experiment after 1 month of in vitro experiment with combined application of three potent PGPR (RG8, RG12 and RGKP3) strain with ZnO NPs. focusing on SSR expression. Duncan's method was used to compare the means at a 1% probability level, with columns sharing the same letters indicating no significant differences	120
Figure 8.3	Estimation of (a) total chlorophyll content; (b) carotenoids content; (c) flavonoids content; (d) proline content for differently treated plants; (e) produced total protein content; and (f) total sugar content in plants with a combination of PGPRs RG8, RG12 and RGKP3 and ZnO NPs in a pot experiment. focusing on SSR expression. Duncan's method was used to compare the means at a 1% probability level, with columns sharing the same letters indicating no significant differences	121

List of Tables

Table No.	Title	Page no.
Table 2.1	Effect of various nanoparticles on plants with PGPR	26
Table 3.1	Qualitative and Quantitative analysis of PGP Traits	36
Table 5.1	Effect of Different PGPRs on Seed Germination and Vigor Index	68

List of Abbreviation

%	:	Percent
°C	:	Degree of Celsius
CO₂	:	Carbon dioxide
PGPR	:	Plant growth promoting rhizobacteria
NPs	:	Nanoparticles
ZnO	:	Zinc oxide
IAA	:	Indole-3-acetic acid
HCN	:	Hydrogen cyanide
OD	:	Optical density
XRD	:	X-ray diffraction
EM	:	Scanning electron microscopy
TEM	:	Transmission electron microscopy
SOFI	:	State of food security
IOPEPC	:	Indian oilseeds and produce export promotion council
GA	:	Gibberellin
ACC	:	1-Aminocyclopropane-1-Carboxylate Deaminase
KSB	:	Potassium solubilizing bacteria
DNA	:	Deoxyribonucleic acid
ROS	:	Reactive oxygen species
ATP	:	Adenosine triphosphate
AgNPs	:	Silver nanoparticles
SiO₂	:	Silicon dioxide
BSA	:	Bovine serum albumin
CaP NPs	:	Calcium phosphate nanoparticles
Mn NPs	:	Manganese nanoparticles.
VOCs	:	Volatile organic compounds
EPS	:	Extracellular polymeric substances,
Fe₃O₄	:	Magnetite nanoparticles
CuO	:	Copper oxide
SSR	:	Regression sum of square
MONPs	:	Molybdenum oxide nanoparticles
